

LBA Calibrator Survey

OPAL LBA Cover Sheet

Name	Email	Affiliation	Country	Student	Available for observations?
Principal investigator					
Leonid Petrov	Leonid.Petrov@lpetrov.net	NVI, Inc./NASA GSFC	United States	No	No
Co-Investigators					
Chris Phillips	Chris.Phillips@csiro.au	Australia Telescope National Facility	Australia	No	Yes
Alessandra Bertarini	abertari@mpifr-bonn.mpg.de	Max Planck Inst. fuer Radioastronomie, Bonn	Germany	No	No
Ed Fomalont	efomalon@nrao.edu	National Radio Astronomy Observatory	United States	No	No
Anastasios Tzioumis	tasso.tzioumis@csiro.au	Australia Telescope National Facility	Australia	No	Yes
Sergei Pogrebenko	pogrebenko@jive.nl	Joint Institute for Very Long Baseline Interferometry in Europe	Netherlands	No	No
Tara Murphy	tara@physics.usyd.edu.au	University of Sydney	Australia	No	No
Elaine Sadler	ems@physics.usyd.edu.au	University of Sydney	Australia	No	No
Sarah Burke	sarahgburke@gmail.com	Swinburne University of Technology	Australia	Yes	No
Kee-Tae Kim	ktkim@kasi.re.kr	Korea Astronomy & Space Science Institute	Korea, Republic of	No	No
Ron Ekers	ron.ekers@csiro.au	Australia Telescope National Facility	Australia	No	No
Roy Booth	roy@hartrao.ac.za	Hartebeesthoek Radio Astronomical Observatory	South Africa	No	No

Proposal Details

Proposal type	Standard
Continuing	No
Scientific categories	Extragalactic
Help required	None

Instrument Information

Antennas requested

- Parkes
- ATCA (single)
- Mopra
- Tidbinbilla (Any)
- Hobart
- Ceduna
- HartRAO

Observing mode Disk recording

Simultaneous ATCA observations No

Recorder LBADR

Other information Astrometry

Correlation location Bonn

Shipping location Bonn, Address: Contents of disks from ATCA, Ceduna, and Mopra will be transferred electronically to Bonn. Disks from Hobart26, Hartrao, Parkes, and Tidbinbilla will be shipped directly to Bonn from the stations.

Abstracts

Scientific

We request two 24 hour observing sessions at ATCA, Ceduna, Mopra, Hobart, Parkes, Hartrao (and Tidbinbilla if available) for running X-band observations to determine coordinates at a milliarcsec accuracy level for ~400 flat-spectrum sources in the declination zone [-45, -90]. We will also produce estimates of correlated flux density and source structure. The output catalogue of source positions will be of use for phase-referencing observations at the LBA, as a calibrator pool for the ATCA, ALMA, and SKA, for space navigation, and as a source list for geodetic observations. Considering success rate of 75%, the pool of known southern calibrators is expected to increase by a factor of 4.

Outreach

The goal of our project is to improve the map of southern sky in radiowaves. Positions of 400 compact sources will be determined with the highest accuracy that the modern technology allows. For the first time astronomers will make images of radiogalaxies that are never seen in the northern hemisphere.

Scheduling

Special requirements	24hr blocks
Total time for project	120 hours (previous + this proposal + future requests)
Allocated time so far	0 hours (all previous semesters)

OPAL LBA Observations Table

Name	Position			Integration time (hours)	Repeats	Total time	Target type	Band	Polarisations	IFs	Frequencies (MHz)	Bandwidths (MHz)	Transitions	Data rate (Mbps)
	RA	Dec	Epoch											
400 target sources	00:00:00	-90:00:00	J2000	44.0	1	44.0	many sources	3cm	2	4	8400	16		512
50 tropospheric calibrators	00:00:00	-90:00:00	J2000	4.0	1	4.0	many sources	3cm	2	4	8400	16		512

Total time for semester: 48.0 hours

LBA Calibrator Survey

1 Summary

We request two 24 hour observing sessions at ATCA, Ceduna, Mopra, Hobart, Parkes, Hartrao (and Tidbinbilla if available) for running X-band observations to determine coordinates at a milliarcsec accuracy level for ~ 400 flat-spectrum sources in the declination zone $[-45^\circ, -90^\circ]$. We will also produce estimates of correlated flux density and source structure. The output catalogue of source positions will be of use for phase-referencing observations at the LBA, as a calibrator pool for the ATCA, ALMA, and SKA, for space navigation, and as a source list for geodetic observations. Considering success rate of 75%, the pool of known southern calibrators is expected to increase by a factor of 4.

This project extends the existing approved project v254a (Phillips et al.). The scope and source selection are essentially unchanged, but analysis approach is much improved. As such, this current project should be considered to replace v254a.

2 Why calibrators are needed

A catalogue of compact radio sources with position known at to an accuracy of several milliarcsec is needed for many applications. Among them are phase referencing for imaging weak objects with VLBI, accurate differential astrometry (for example, for pulsar and maser proper motions and parallax), spacecraft navigation, calibrators for ALMA, targets for space navigation, monitoring the Earth's rotation, and space geodesy.

3 Why new observations are needed

To date, positions of 3913 sources¹ were produced from analysis of more than 4400 24 hour VLBI experiments from 1979 through present, and from 27 VLBA Calibrator Survey experiments that the team of proposers scheduled and analyzed [1, 2, 3, 4, 5]. However, the vast majority of known calibrators lie in the northern hemisphere accessible to the VLBA. Plot 1 shows the sky calibrator density in 4° declination zones as a function of the zone declination. We see the rapid drop of source calibrator density starting with declinations -30° and further south. There are only 98 calibrators in the declination zone $[-90^\circ, -52^\circ]$ versus 524 objects in the declination zone $[+52^\circ, +90^\circ]$. Positions of known 98 calibrators in the $[-90^\circ, -52^\circ]$ zone were derived from geodetic observations at a single baseline Hartrao/Hobart over 15 years. The probability to find a calibrator within a circle of 4° , 3° , and 2° at any given direction in various declination zones is shown in table 1. To reach such a calibrator density that within 3° at any given direction in the declination zone $[-45^\circ, -90^\circ]$ at least one calibrator will be found with the probability $> 90\%$, in total, ~ 1000 objects have to be observed. In this proposal we will concentrate on ~ 400 brightest objects, and we will request time to observe further sources later.


In order to address the problem of poor calibrator coverage of the southern hemisphere, we propose two 24 hour observing sessions with the LBA. We propose to observe only those sources that the VLBA cannot reach, i.e. sources in the declination zone $[-90^\circ, -45^\circ]$. We have plans to submit a separate proposal to the VLBA and to find more calibrators in the declination zone $[-52^\circ, -30^\circ]$.

¹<http://vlbi.gsfc.nasa.gov/astro>

Table 1: Probability to find a calibrator in the disk of the specified radius at any given direction in various declination zones

Zone	2°	3°	4°
-20°, +90°	70.2%	94.1%	99.5%
-30°, -20°	64.6%	90.7%	98.5%
-40°, -30°	45.1%	75.1%	91.6%
-52°, -40°	21.5%	42.3%	61.2%
-90°, -52°	21.3%	41.4%	61.1%

Figure 1: Calibrator source density per steradian as a function in various declination zones



4 Past observations

On 2007.06.24 a 12 hour geodetic experiment v230, using ATCA, Mopra, Ceduna, Hobart, Parkes, Seshan, and Hartrao was observed. The goal of that experiment was to determine accurate positions of ATCA, Mopra, Ceduna with centimeter level of accuracy. In addition to geodetic sources, five new sources were scheduled in this experiment. As of time of writing, 2007.12.12_21:00 UTC, the experiment is being correlated in Bonn. Fringes have been found at all 28 baselines². including baselines recorded with LBADR and Mark5. Using 8 hours of the data at 6 baselines that have been correlated (17% of the experiment), we have run a preliminary solution. We found that a) all group delay ambiguities with spacings 4.8077 ns were successful resolved, no outliers found, all points were used in the solution; b) position of Ceduna and Mopra were determined with precision of 10–12 cm for the vertical component and 2–3 cm for the horizontal component; c) position accuracy of two new sources J0220-6330 and J1945-5520 detected in one scan at two baselines, Ceduna/Mopra and Mopra/Hobart, was 10–20 mas. Including in analysis all 20 scheduled observations at 10 Australian baselines, will improve the position accuracy by a factor of $\sqrt{10}$, and including 9000 km long baselines with Hartrao will improve the accuracy even further.

We consider that the previous experiment has proved the concept of the proposed observations.

²See <http://vlbi.gsfc.nasa.gov/pet/discussion/v230/> for the current status of v230 analysis.

5 Proposed observations

We propose to observe two 24 hours sessions at X-band at ATCA, Ceduna, Mopra, Hobart, Parkes, and Hartrao. All stations will be observing at 8.4 GHz at four IF channels of 16 MHz each spread over 256 MHz with two polarizations. The stations ATCA, Ceduna, Mopra will be utilizing the LBADR disk based recording system, while Parkes, Hobart, Seshan25, Hartrao will use Mark5. The data will be correlated on the geodetic correlator at Bonn. Data from ATCA, Ceduna, Mopra and Parkes will be transferred to Bonn via the network, where they will be re-written into Mark5b. This was also successfully demonstrated for v230.

We request any Tidbinbilla antenna, if available, for the improved uv -coverage. While the observations do not depend on Tidbinbilla, the extra baselines will improve the accuracy of the solutions. Results of these observations will be of interest for JPL for support of future interplanetary missions and for support of Plank mission.

Candidate flat-spectrum sources in the declination range of $[-90^\circ, -45^\circ]$ from the AT20G catalogue will be selected on the basis of their probability of being detected using the algorithm developed and successfully tested in [4]. The catalogue AT20G will provide input for that algorithm: estimates of flux densities at 8.4 GHz and spectral indexes.

We will apply the same scheduling strategy that we have successfully applied for the VLBA Calibrator Survey [2]. The sources will be observed in a sequence that minimizes slewing in two scans of 120 seconds each. Every 1.5 hours a block of 3–4 strong sources with precisely known positions and recent maps, so called tropospheric calibrators, will be observed. Tropospheric calibrators will be scheduled in such a manner that each station will observe at least one source at the elevation range of $[5^\circ, 20^\circ]$ (except Parkes), one source at the elevation range of $[20^\circ, 50^\circ]$, and one source at the elevation range of $[50^\circ, 90^\circ]$. The tropospheric calibrators will also serve two other goals: 1) they will tie the resulting catalogue to the ICRF catalogue; 2) they will be used as additional amplitude calibrators.

Final data analysis will be performed at NASA using Calc/Solve software program in a similar way how positions of other 3913 sources have been determined. The expected accuracy of source positions is 1–5 mas. We will be using global GPS maps for alleviation of ionospheric errors. According to our experience of processing VLBA X-band data, unaccounted ionosphere errors will set the accuracy limit of 0.5–2.0 mas depending on the ionosphere condition.

6 Outcomes

We will publish on the web positions from each epoch within 21 days upon completion of correlation. Once the full survey is complete, we will publish the full source list with flux densities and positions and make a web searchable interface to the list.

These observations will also be used for further improving site coordinates.

The full survey will certainly also spur numerous science projects, such as investigating the AT20G population or searching for specific objects such as gravitational lenses. One such example is proposal v275, a search for super-massive black holes also being submitted this term.

References

- [1] Beasley, A. J. et al., ApJS, 141, 13.
- [2] Fomalont, E., Petrov, L., McMillan, D. S., Gordon, D., Ma, C. 2003, AJ, 126, 2562.
- [3] Petrov, L., Kovalev, Y.Y., Fomalont, E., Gordon, D., 2005, AJ, 129, 1163.
- [4] Petrov, L., Yu.Y. Kovalev, E. Fomalont, D. Gordon, 2006, AJ, 131, 1872.
- [5] Kovalev, Yu.Y., L. Petrov, E. Fomalont, D. Gordon, 2007, AJ, 133, 1236.